

# ADISSEO: UNITED STATES OF AMERICA 2019 MYCOTOXIN SURVEY IN CORN AND CORN SILAGE

Author: Radka Borutova DVM, PhD; Scientific & Technical Manager at ADISSEO France

## Introduction

The presence of mycotoxins in feed and foodstuffs is an important concern for human and animal health. The *Aspergillus*, *Fusarium* and *Claviceps* species of fungi are ubiquitous in nature and under ideal conditions, they often can infect crops and forages in fields, during storage, shipment and processing. Mycotoxins can cause serious health problems in animals and their presence in agricultural commodities may result in serious economic losses. Mycotoxins have been implicated in a variety of human diseases, including some forms of cancer. It has been estimated that mycotoxin-contaminated grains cost grain handlers and the livestock industry several hundred million dollars annually (CAST, 2003).



Picture 1: Corn harvested in Iowa (US) 2019  
©Copyright: Radka Borutova, Adisseo France

The 2019 Adisseo mycotoxin survey included 48 corn and silage samples from across the United States of America (USA) (i.e. Iowa, Illinois, Nebraska, Minnesota, Missouri, South Dakota, Pennsylvania, New York, Wisconsin, Idaho, Ohio). All samples were collected almost immediately after the harvest from farms or animal feed production sites. Sample providers were advised to follow the principles of good sampling (Richard, 2000). Analytical personnel and/or laboratory staff were not involved in the sampling and did not influence any part of this procedure.

A total of 48 analyses were conducted to test for the occurrence of the 11 mycotoxins most commonly found in agricultural commodities intended for animal production. The survey provided an insight into the incidences of aflatoxin B<sub>1</sub> (Afb<sub>1</sub>), aflatoxin B<sub>2</sub> (Afb<sub>2</sub>), aflatoxin G<sub>1</sub> (Afg<sub>1</sub>), aflatoxin G<sub>2</sub> (Afg<sub>2</sub>), ochratoxin A (OTA), zearalenone (ZEN), deoxynivalenol (DON), T-2 toxin, HT-2 toxin, fumonisin B<sub>1</sub> (FB<sub>1</sub>) and fumonisin B<sub>2</sub> (FB<sub>2</sub>).

All 11 mycotoxins were analyzed by liquid chromatography tandem mass spectrometry (LC MS/MS). For the purpose of data analysis, non-detection levels were based on the limits of quantification (LOQ) of the test method for each mycotoxin: Afb<sub>1</sub> <20 µg/kg; Afb<sub>2</sub> <20 µg/kg; Afg<sub>1</sub> <20 µg/kg; Afg<sub>2</sub> <20 µg/kg; ZEN <100 µg/kg; DON <200 µg/kg; FB<sub>1</sub> <200 µg/kg; FB<sub>2</sub> <200 µg/kg; OTA <20 µg/kg; T-2 toxin <20 µg/kg and HT2-toxin <200 µg/kg.

## Results

The results showed that 4.5% of the silage samples and 34.6% of the corn samples were contaminated with ZEN (Table 1), a mycotoxin which can affect the fertility performance of most species. The average concentration of ZEN in silage was 158 µg/kg and 527.90 µg/kg in corn. These levels of ZEN in corn are considered to carry a medium risk and can be harmful especially to dairy cows and herd bulls. In ruminants such as cows and bulls, ZEN is converted by the rumen flora into its major component, hydroxy-metabolite  $\alpha$ -zearalenol (approximately 90%) and to a lesser extent to  $\beta$ -zearalenol (Kennedy et al., 1998).  $\alpha$ -zearalenol has a higher estrogenic potency compared to the parent ZEN (Seeling et al., 2005).


86.4% of the silage samples and all of the corn samples were contaminated with DON. There was a high average concentration of DON of 1370.6 µg/kg and a maximum concentration of 4357 µg/kg detected in one of the silage samples. The highest concentration of DON detected in one of the corn samples was almost the same as the concentration detected in silage and was equally high at 4118 µg/kg. A medium average concentration of 966.4 µg/kg of DON was detected in corn.

As expected, 50% of corn and 18.2 % of silage samples contained FB<sub>1</sub>. The maximum concentration of FB<sub>1</sub> toxin found in one of the corn samples was 4574 µg/kg. This high concentration is usual for corn grown in the USA and may have a significant effect on the health and performance of farm animals, especially swine and horses.

T-2 toxin was detected in 15.4% of corn samples and 4.5% of silage samples. OTA, Afb<sub>1</sub>, Afb<sub>2</sub>, Afg<sub>1</sub>, Afg<sub>2</sub> and HT-2 toxins were not detected in any of the analyzed samples.

Several samples had 2 to 4 different mycotoxins detected which could lead to synergistic interactions among them. In total, 22%, 15% and 11% of samples were contaminated with 2, 3 and 4 mycotoxins, respectively (Figure 1).

The average concentrations of all the detected mycotoxins were low to medium (>LOQ and above US Food and Drug Administration [FDA] regulatory and advisory levels).




**Table 1:** Mycotoxin contamination of corn in the USA 2019

Mycotoxins	% of positive samples		Average concentration of positive samples (µg/kg)		Maximum concentration of positive mycotoxin (µg/kg)	
	Corn	Silage	Corn	Silage	Corn	Silage
AfB <sub>1</sub>	ND	ND	ND	ND	ND	ND
AfB <sub>2</sub>	ND	ND	ND	ND	ND	ND
AfG <sub>1</sub>	ND	ND	ND	ND	ND	ND
AfG <sub>2</sub>	ND	ND	ND	ND	ND	ND
DON	100.0	86.4	966.4	1370.6	4118.0	4357.0
ZEN	34.6	4.5	527.9	158.0	2022.0	158.0
FB <sub>1</sub>	50.0	18.2	1771.8	424.3	4574.0	385.0
FB <sub>2</sub>	30.0	ND	711.3	ND	1423.0	ND
T2	15.4	4.5	29.5	28.0	33.0	28.0
HT-2	ND	ND	ND	ND	ND	ND
OTA	ND	ND	ND	ND	ND	ND

AfB<sub>1</sub> = aflatoxin B<sub>1</sub>  
 AfB<sub>2</sub> = aflatoxin B<sub>2</sub>  
 AfG<sub>1</sub> = aflatoxin G<sub>1</sub>  
 AfG<sub>2</sub> = aflatoxin G<sub>2</sub>  
 DON = deoxynivalenol  
 FB<sub>1</sub> = fumonisin B<sub>1</sub>  
 FB<sub>2</sub> = fumonisin B<sub>2</sub>  
 OTA = ochratoxin A  
 ZEN = zearalenone  
 T2 = T2 toxin  
 HT2 = HT<sub>2</sub> toxin  
 ND = not detected

**Average number of mycotoxins per sample**



**Figure 1:** Average number of mycotoxins per sample [%]

## Conclusion

The Adisseo 2019 mycotoxin survey concluded that the 2019 corn and corn silage harvest in the USA was of medium (>LOQ but below FDA regulatory and advisory levels) quality in terms of mycotoxin contamination.

Based on the results of this survey which was conducted immediately after the 2019 harvest, the 2019 corn crop in the USA should not automatically be considered safe for inclusion into finished feed rations for all animal species. A degree of vigilance is prudent.

The last possible line of defense is the application of effective mycotoxin management strategies. The application of mycotoxin management strategy is a very common method to prevent mycotoxicosis and is an effective strategy to keep mycotoxin risk low under any and all conditions

## References

- Kennedy DG, Hewitt SA, McEvoy JD, Currie JW, Cannavan A, Blanchflower WJ, Elliot CT. 1998. Zeranol is formed from Fusarium spp. toxins in cattle in vivo. Food Addit Contam 15:393-400.
- Seeling K, Dänicke S, Ueberschar KH, Lebzien P, Flachowsky G. 2005. On the effects of Fusarium toxin contaminated wheat and the feed intake level on the metabolism and carry over of zearalenone in dairy cows. Food Addit Contam 22:847-855.
- Richard, J., 2000. Sampling and sample preparation for mycotoxin analysis. Romer® Labs Guide to Mycotoxins. 2. Romer® Labs Inc., 1301 Stylemaster Drive, Union, MO, USA 63084-1156.
- FDA Mycotoxin Regulatory Guidance, 2011, A Guide for Grain Elevators, Feed Manufacturers, Grain Processors and Exporters, National Grain and Feed Association, 1250 Eye St., N.W., Suite 1003, Washington, D.C., 20005-3922.
- CAST, 2003. Mycotoxins: Risks in Plant, Animal and Human Systems. Task Force Report No.139. Council for Agricultural Science and Technology. Ames, IA.

Adisseo is one of the world's leading experts in feed additives. The group has 10 research centers and production sites in Europe, USA and China to design, produce and market nutritional solutions for sustainable animal feed. With more than 2,200 employees, Adisseo serves around 3,900 customers in over 110 different countries through its global distribution network. Adisseo is one of the main subsidiaries of China National BlueStar, a leader in the Chinese chemical industry with nearly 21,500 employees and a turnover of 9,3 billion USD. Adisseo is listed on the Shanghai Stock Exchange. Adisseo is listed on the Shanghai Stock Exchange. **Find out more at [www.adisseo.com](http://www.adisseo.com)**

